arXiv:1303.1582v1 [math.CA] 7 Mar 2013

# COMPLETE MONOTONICITY OF A DIFFERENCE BETWEEN THE EXPONENTIAL AND TRIGAMMA FUNCTIONS

FENG QI AND XIAO-JING ZHANG

Abstract. In the paper, by directly verifying an inequality which gives a lower bound for the first order modified Bessel function of the first kind, the authors supply a new proof for the complete monotonicity of a difference between the exponential function $e^{1/t}$ and the trigamma function $\psi'(t)$ on $(0,\infty)$.

## 1. Introduction

In [3, Lemma 2], the inequality

$$\psi'(t) < e^{1/t} - 1 \tag{1.1}$$

on $(0,\infty)$ was discovered and employed, where $\psi(t)$ denotes the digamma function

$$\psi(t) = [\ln\Gamma(t)]' = \frac{\Gamma'(t)}{\Gamma(t)}$$

and $\Gamma$ is the classical Euler gamma function which may be defined for $\Re(z) > 0$ by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} \,\mathrm{d}\, t.$$

The functions $\psi'(z)$ and $\psi''(z)$ are respectively called the trigamma function and the tetragamma function. As a whole, the derivatives $\psi^{(k)}(z)$ for $k \in \{0\} \cup \mathbb{N}$ are called polygamma functions.

An infinitely differentiable function $f$ defined on an interval $I$ is said to be a completely monotonic function on $I$ if it satisfies

$$(-1)^k f^{(k)}(x) \ge 0 \tag{1.2}$$

for all $k \in \{0\} \cup \mathbb{N}$ on $I$. Some properties of the completely monotonic functions can be found in, for example, [2, 7].

In [5, Theorem 3.1] and [6, Theorem 1.1], the following theorem was proved totally by three methods.

**Theorem 1.1.** *The function*

$$h(t) = e^{1/t} - \psi'(t) \tag{1.3}$$

*is completely monotonic on* $(0,\infty)$ *and*

$$\lim_{t\to\infty} h(t) = 1. \tag{1.4}$$

---

2010 *Mathematics Subject Classification.* Primary 26A48, 33B10, 33C10; Secondary 26A12, 26D07, 33B15, 33C20, 44A10.

*Key words and phrases.* Complete monotonicity; completely monotonic function; integral representation; difference; exponential function; trigamma function; inequality; modified Bessel function of the first kind.

The second main result of the paper [6] is its Theorem 1.2 which has been referenced in [4, Section 1.2] and [5, Lemma 2.1] as follows.

**Theorem 1.2.** *For $k \in \{0\} \cup \mathbb{N}$ and $z \ne 0$, let*

$$H_k(z) = e^{1/z} - \sum_{m=0}^{k} \frac{1}{m!} \frac{1}{z^m}. \tag{1.5}$$

*For $\Re(z) > 0$, the function $H_k(z)$ has the integral representations*

$$H_k(z) = \frac{1}{k!(k+1)!} \int_0^\infty {}_1F_2(1; k+1, k+2; t) t^k e^{-zt} \,\mathrm{d} t \tag{1.6}$$

*and*

$$H_k(z) = \frac{1}{z^{k+1}} \biggl[\frac{1}{(k+1)!} + \int_0^\infty \frac{I_{k+2}\bigl(2\sqrt{t}\,\bigr)}{t^{(k+2)/2}} e^{-zt} \,\mathrm{d} t\biggr], \tag{1.7}$$

*where the hypergeometric series*

$${}_pF_q(a_1, \dots, a_p; b_1, \dots, b_q; x) = \sum_{n=0}^\infty \frac{(a_1)_n \cdots (a_p)_n}{(b_1)_n \cdots (b_q)_n} \frac{x^n}{n!} \tag{1.8}$$

*for $b_i \notin \{0, -1, -2, \dots\}$, the shifted factorial $(a)_0 = 1$ and*

$$(a)_n = a(a+1) \cdots (a+n-1) \tag{1.9}$$

*for $n > 0$ and any real or complex number $a$, and the modified Bessel function of the first kind*

$$I_\nu(z) = \sum_{k=0}^\infty \frac{1}{k!\Gamma(\nu+k+1)} \left(\frac{z}{2}\right)^{2k+\nu} \tag{1.10}$$

*for $\nu \in \mathbb{R}$ and $z \in \mathbb{C}$.*

When $k = 0$, the integral representations (1.6) and (1.7) may be written as

$$e^{1/z} = 1 + \int_0^\infty \frac{I_1\bigl(2\sqrt{t}\,\bigr)}{\sqrt{t}} e^{-zt} \,\mathrm{d} t \tag{1.11}$$

and

$$e^{1/z} = 1 + \frac{1}{z} \biggl[1 + \int_0^\infty \frac{I_2\bigl(2\sqrt{t}\,\bigr)}{t} e^{-zt} \,\mathrm{d} t\biggr] \tag{1.12}$$

for $\Re(z) > 0$. Hence, by the well known formula

$$\psi^{(n)}(z) = (-1)^{n+1} \int_0^\infty \frac{u^n}{1 - e^{-u}} e^{-zu} \,\mathrm{d} u \tag{1.13}$$

for $\Re(z) > 0$ and $n \in \mathbb{N}$, see [1, p. 260, 6.4.1], the function $h(t)$ defined by (1.3) has the following integral representation

$$h(t) = 1 + \int_0^\infty \biggl[\frac{I_1\bigl(2\sqrt{u}\,\bigr)}{\sqrt{u}} - \frac{u}{1 - e^{-u}}\biggr] e^{-tu} \,\mathrm{d} u. \tag{1.14}$$

**Proposition 1.1** (Hausdorff-Bernstein-Widder Theorem [7, p. 161, Theorem 12b])**.** *A necessary and sufficient condition that $f(x)$ should be completely monotonic for $0 < x < \infty$ is that*

$$f(x) = \int_0^\infty e^{-xt} \,\mathrm{d} \alpha(t), \tag{1.15}$$

*where $\alpha(t)$ is non-decreasing and the integral converges for $0 < x < \infty$.*

Combining the complete monotonicity in Theorem 1.1 and the integral representation (1.14) with the necessary and sufficient condition in Proposition 1.1, it was revealed in [6] that

$$\frac{I_1\bigl(2\sqrt{u}\,\bigr)}{\sqrt{u}} \ge \frac{u}{1-e^{-u}}, \quad u>0. \tag{1.16}$$

Replacing $2\sqrt{u}$ by $t$ in (1.16) yields [6, Theorem 1.3] below.

**Theorem 1.3.** *For $t>0$, we have*

$$I_1(t) > \frac{(t/2)^3}{1-e^{-(t/2)^2}}. \tag{1.17}$$

We note that the complete monotonicity in Theorem 1.1 is the basis of the inequality (1.17) and some results in the subsequent papers [4, 5].

The aim of this paper is, with the help of the integral representation (1.14) but without using Proposition 1.1, to supply a new proof of Theorems 1.1 and 1.3 in a converse direction with that in [4, 5, 6]. In other words, Theorem 1.3 will be firstly and straightforwardly proved, and then Theorem 1.1 will be done.

## 2. A NEW PROOF OF THEOREMS 1.1 AND 1.3

By the definition of the modified Bessel function $I_\nu(z)$ in (1.10), it is easy to see that

$$\frac{I_1\bigl(2\sqrt{u}\,\bigr)}{\sqrt{u}} = \sum_{k=0}^{\infty}\frac{1}{k!\Gamma(k+2)}u^k > 1+\frac{1}{2}u+\frac{1}{12}u^2.$$

Hence, in order to prove (1.16), it suffices to show

$$1+\frac{1}{2}u+\frac{1}{12}u^2 \ge \frac{u}{1-e^{-u}} \tag{2.1}$$

which is equivalent to

$$\begin{aligned}
&e^u\bigl(12-6u+u^2\bigr)-12-6u-u^2\\
&> \biggl(1+u+\frac{u^2}{2}+\frac{u^3}{3!}+\frac{u^4}{4!}+\frac{u^5}{5!}\biggr)\bigl[3+(u-3)^2\bigr]-12-6u-u^2\\
&= \frac{1}{120}u^5\biggl[\frac{3}{4}+\biggl(\frac{1}{2}-u\biggr)^2\biggr]\\
&\ge 0.
\end{aligned}$$

Consequently, the proof of the inequality (1.16), that is, Theorem 1.3, is thus complete.

Substituting the inequality (1.16) into the integral representation (1.14) leads to $h(t)>0$ and for $k\in\mathbb{N}$

$$(-1)^k h^{(k)}(t) = \int_0^\infty \biggl[\frac{I_1\bigl(2\sqrt{u}\,\bigr)}{\sqrt{u}} - \frac{u}{1-e^{-u}}\biggr] u^k e^{-tu}\,\mathrm{d}\,u > 0$$

on $(0,\infty)$. As a result, the function $h(t)$ is completely monotonic on $(0,\infty)$.

The limit (1.4) follows immediately from taking $t\to\infty$ on both sides of the integral representation (1.14). Theorem 1.1 is proved.

*Remark* 2.1. The inequality (2.1) is equivalent to

$$Q(u)=e^u\big(12-6u+u^2\big)-12-6u-u^2>0.$$

An immediate differentiation yields

$$\begin{aligned}Q'(u)&=e^u\big(u^2-4u+6\big)-2(u+3),\\ Q''(u)&=e^u\big(u^2-2u+2\big)-2,\\ Q'''(u)&=u^2e^u.\end{aligned}$$

Since $Q'''(u)$ and $Q''(0)=0$, it follows that $Q''(u)>0$ on $(0,\infty)$. Owing to $Q'(0)=0$ and $Q''(u)>0$, it is derived that $Q'(u)>0$. Finally, since $Q(0)=0$, the function $Q(u)$ is positive on $(0,\infty)$. This gives an alternative proof of the inequality (2.1).

## References

[1] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, 9th printing, Washington, 1970.

[2] B.-N. Guo and F. Qi, *A completely monotonic function involving the tri-gamma function and with degree one*, Appl. Math. Comput. **218** (2012), no. 19, 9890–9897; Available online at `http://dx.doi.org/10.1016/j.amc.2012.03.075`.

[3] B.-N. Guo and F. Qi, *Refinements of lower bounds for polygamma functions*, Proc. Amer. Math. Soc. **141** (2013), no. 3, 1007–1015; Available online at `http://dx.doi.org/10.1090/S0002-9939-2012-11387-5`.

[4] F. Qi, *Properties of modified Bessel functions and completely monotonic degrees of differences between exponential and trigamma functions*, available online at `http://arxiv.org/abs/1302.6731`.

[5] F. Qi and C. Berg, *Complete monotonicity of a difference between the exponential and trigamma functions and properties related to a modified Bessel function*, Mediterr. J. Math. **10** (2013), in press; Available online at `http://dx.doi.org/10.1007/s00009-013-0272-2`.

[6] F. Qi and S.-H. Wang, *Complete monotonicity, completely monotonic degree, integral representations, and an inequality related to the exponential, trigamma, and modified Bessel functions*, available online at `http://arxiv.org/abs/1210.2012`.

[7] D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1946.

(Feng Qi) Department of Mathematics, School of Science, Tianjin Polytechnic University, Tianjin City, 300387, China; School of Mathematics and Informatics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China

*E-mail address*: `qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com`

*URL*: `http://qifeng618.wordpress.com`

(Xiao-Jing Zhang) Department of Mathematics, School of Science, Tianjin Polytechnic University, Tianjin City, 300387, China

*E-mail address*: `xiao.jing.zhang@qq.com`